


# Hip Sprain: Care Instructions



© Healthwise, Incorporated

## Overview

---

A hip sprain occurs when you stretch or tear ligaments around your hip. Ligaments are tough tissues that connect one bone to another. You can injure your hip in a fall, when you run, or during sports that involve twisting or sudden direction changes, such as basketball or soccer.

Most minor hip sprains get better with treatment at home.

**Follow-up care is a key part of your treatment and safety.** Be sure to make and go to all appointments, and call your doctor if you are having problems. It's also a good idea to know your test results and keep a list of the medicines you take.

## How can you care for yourself at home?

---

- If your doctor gave you crutches or a walker, use them as directed.
- Rest and protect your hip. Try to stop or reduce any action that causes pain.
- Put ice or a cold pack on your hip for 10 to 20 minutes at a time. Try to do this every 1 to 2 hours for the next 3 days (when you are awake) or until the swelling goes down. Put a thin cloth between the ice and your skin.
- Be safe with medicines. Read and follow all instructions on the label.
  - If the doctor gave you a prescription medicine for pain, take it as prescribed.
  - If you are not taking a prescription pain medicine, ask your doctor if you can take an over-the-counter medicine.
- For the first day or two after an injury, avoid things that might increase swelling, such as hot showers, hot tubs, or hot packs.
- After 2 to 3 days, put a heating pad (set on low) or warm moist cloth on your hip before you do light stretches.
- Do exercises to make your hip stronger, as directed by your doctor or physical therapist.

- Return to your usual level of activity as your hip gets better.

## When should you call for help?

---



**Call your doctor now** or seek immediate medical care if:

- Your pain is worse.
- You cannot walk or stand without help.
- You have signs of infection, such as a fever or increased pain, swelling, redness, or warmth in your hip.
- You have signs of a blood clot, such as:
  - Pain in your calf, back of the knee, thigh, or groin.
  - Redness and swelling in your leg or groin.
- You have tingling, weakness, or numbness in your leg, foot, or toes.

Watch closely for changes in your health, and be sure to contact your doctor if:

- Your pain does not get better in 2 or 3 days.
- You still have pain after 2 weeks.

### Where can you learn more?

---

Go to <https://www.healthwise.net/patiented>

Enter **G946** in the search box to learn more about "**Hip Sprain: Care Instructions**".

---

©2006-2022 Healthwise, Incorporated.

This care instruction is for use with your licensed healthcare professional. If you have questions about a medical condition or this instruction, always ask your healthcare professional. Healthwise, Incorporated disclaims any warranty or liability for your use of this information.